

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 738 706 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
23.10.1996 Bulletin 1996/43

(51) Int. Cl.⁶: C07C 229/42, C07C 235/08,
A61K 31/215, A61K 31/165

(21) Application number: 96106009.2

(22) Date of filing: 17.04.1996

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE

(30) Priority: 19.04.1995 ES 9500756


(71) Applicant: PRODES, S.A.
08960 Sant Just Desvern, Barcelona (ES)

(72) Inventors:
• Serra Masia, Xavier
08920 Cerdanyola del Valles, Barcelona (ES)
• Pi Sallent, Joan
08120 La Llagosta, Barcelona (ES)

(74) Representative: Modiano, Guido, Dr.-Ing. et al
Modiano, Josif, Pisanty & Staub,
Baaderstrasse 3
80469 München (DE)

(54) Nitric esters of 2-(2,6-dihalophenylamino) phenylacetoxyacetic acid derivatives and their preparation process

(57) Nitric esters of 2-(2,6-dihalophenylamino) phenylacetoxyacetic acid derivatives with the formula (I), where: [A = F, Cl or Br; X = O, NH or NR (R = C₁ - C₈ alkyl); R₁ and R₂ independently = C₁ - C₈ alkyl and n is a whole number from 1 to 10. The procedure includes the condensation of 2-(2,6-dihalophenylamino) phenylacetoxyacetic acid with a compound with the formula:



where Y = OH, NH₂ or NHR and Z is Cl, Br or ONO₂.

EP 0 738 706 A1

Description

This invention refers to the nitric esters of 2-(2,6-dihalophenylamino) phenylacetoxycetic acid derivatives; the invention also refers to the procedures for their preparation.

State of the art

An already classical product of the non-steroid anti-inflammatory group is the sodium salt of the 2-(2,6-dichlorophenylamino) phenylacetic acid. However, it is well known that its use causes severe gastrointestinal and hepatic secondary effects. With the aim of improving this limitation, certain functionalized derivatives with an ester of the nitric acid have been claimed recently (WO 94/04484). These derivatives have been described in the above mentioned patent as having an anti-inflammatory activity with less gastric secondary effects.

Aim of the invention

The actual petitioner owns the patent EP 0 119 932 referred, an AINE, the 2-(2,6-dichlorophenylamino) phenylacetoxycetic acid (Aceclofenaco).


This product, besides having a powerful anti-inflammatory and analgesic activity, presents a very improved gastric tolerance (Drugs and Inflammation, Vol. 32 (1991)) well proved in clinical studies (Clin. Tri. J., 27(1), 12-19 (1990), Cli. Tri. J., 25(2), 144-151 (1988), Curr. Ther. Res., 44(2), 252-256 (1988), Drugs Exp. Clin. Res., 15(1), 47-51 (1989). Due to these qualities, the Aceclofenaco is actually a well known prestigious AINE.

Researches on products related with Aceclofenaco have been carried out with the aim of providing new products that, keeping the advantageous pharmacological qualities of the 2-(2,6-dichlorophenylamino) phenylacetoxycetic acid, allow the complete elimination of the rest of gastric injuries.

The preparation process of the said derivatives of the 2-(2,6-dichlorophenylamino) phenylacetoxycetic acid are also a target of this invention.

Description of the invention

The purpose mentioned in the previous paragraph is obtained by the products of this invention, nitric esters of the 2-(2,6-dihalophenylamino) phenylacetoxycetic acid derivatives represented by the general formula I:




(I)

where:


- A is fluorine, chlorine or bromine;
- X means oxygen, NH or NR where R means a linear or ramified alkyl chain of 1 to 6 carbon atoms;
- R₁ and R₂ mean, independently, hydrogen or a linear or ramified alkyl chain of 1 to 6 carbon atoms; and
- n is a number between 1 to 10.

Among the compounds of this invention, especially interesting are some nitrate compounds of general structure I where A is chlorine, X is oxygen, R₁ and R₂ are both hydrogen and n is between 2 to 4, which are represented by the

following formula II:




Similarly, it is also especially interesting the nitroester with the general formula I, where A is chlorine, X is NH, R₁ and R₂ are both hydrogen and n = 2, which is represented by formula III:




To prepare the nitroesters with the general formula I several procedures can be used, as described below.

a. The first process consists of carrying out the condensation of the corresponding 2-(2,6-dihalophenylamino) phenylacetoxymethyl acid with the formula IV:



where: A is fluorine, chlorine or bromine, with a derivate with the general formula V:




(V)

where Y can be OH, NH₂ or NHR being R a linear or ramified alkyl chain of 1 to 6 carbon atoms, while R₁, R₂ and n have the same meaning as described before for the general formula I.

The reaction is carried out by means of a suitable condensation agent like N,N'-dicyclohexylcarbodiimide or N,N'-carbonyl diimidazole in an aprotic solvent generally of the chlorinated type (chloroform, dichloromethane), ether (tetrahydrofurane) or similar.


b. A second process of preparation comprises the following steps:

i) Condensation of the 2-(2,6-dihalophenylamino) phenylacetoxyacetic acid with the formula IV with a compound with the general formula VI:



(VI)

where Y, n, R₁ and R₂ have the same meaning as for the compound with the formula V and Z is a Cl or Br atom, to provide the compound with the formula VII.




(VII)

where A, X, R₁, R₂, n and Z have the meaning mentioned above.

The reaction is carried out, as mentioned in section i), by means of N,N'-dicyclohexylcarbodiimide or N,N'-carbonyl diimidazole as a condensation agent in the same conditions.

ii) Optional substitution of Z of the compound with the formula VII with a iodine atom through the reaction with sodium iodine to provide the compound with the formula VIII.



where A, X, R₁, R₂ and n have the same meaning as the one given to the compound with the general formula I.

iii) Displacement shift of Z of the compound with the formula VII or, otherwise, of the iodine atom of the compound with the formula VIII with AgNO₃ in acetonitrile as solvent, to provide the corresponding nitroesters with the formula I.

The products with the formula I in this invention are obtained easily according to what has been said in the former description with a series of procedures with few stages which allow a preparation with a good yield and purely. All this is shown in the following examples:


EXAMPLE 1: Preparation of 2-nitrooxyethyl 2-(2,6-dichlorophenylamino) phenylacetoxycetate.

Solve 6.28 g (0.040 mol) of N,N'-carbonyl diimidazole in 150 ml of anhydrous dichloromethane under a dry nitrogen atmosphere. Shake and add for five minutes 13.87 g (0.039 mol) of 2-(2,6-dichlorophenylamino) phenylacetoxycetic acid in four shares.

Shake for five minutes and add a solution of 4.19 g (0.039 mol) of 2-nitrooxyethanol to 50 ml of dichloromethane. Protect from the light and shake for 18 hours. Eliminate the solvent under reduced pressure and solve again in 150 ml of ethyl acetate.

Wash three times with 100 ml of HCl 0.5N, twice with 100 ml of NaHCO₃ at 10% and neutralize by washing twice with a saturated solution of NaCl. Dry on anhydrous sodium sulphate. Filter, eliminate the solvent at a reduced pressure and purify the oil obtained by chromatography in silica gel column.

Elute with dichloromethane and recover 11.92 g (69%) of 2-nitrooxyethyl 2-(2,6-dichlorophenylamino) phenylacetoxycetate in the form of oil that crystallizes (m.p. 61-64 °C) and that has the following spectroscopic features:



IR (KBr, cm^{-1}): 3360 (N-H), 1744 (C=O), 1640 (ONO_2)

^1H -RMN (DCCl_3 , δ): 3.95 (s, 2H, CH_2 (8)), 4.41 (m, 2H, CH_2 (11)), 4.59 (m, 2H, CH_2 (10)), 4.72 (s, 2H, CH_2 (9)), 6.57 (d, 1H, $J_{4,5} = 7.8$ Hz, C-H (4)), 6.69 (s, 1H, N-H), 6.98 (m, 1H, $J_{6,7} = J_{6,5} = 7.8$ Hz, $J_{6,4} = 0.9$ Hz, C-H (6)), 6.99 (t, 1H, $J_{1,2,3} = 8.1$ Hz, C-H (1)), 7.15 (m, 1H, $J_{5,4} = J_{5,6} = 7.8$ Hz, $J_{5,7} = 1.8$ Hz, C-H (5)), 7.27 (m, 1H, $J_{7,6} = 7.8$ Hz, $J_{7,5} = 1.8$ Hz, C-H (7)), 7.35 (d, 2H, $J_{2,3,1} = 8.1$ Hz, C-H (2)(3)).

^{13}C -RMN (DCCl_3 , δ): 37.94, 60.93, 61.01, 69.82, 118.42, 122.18, 123.71, 124.13, 128.21, 128.86, 129.49, 130.96, 137.73, 142.67, 167.13, 171.38.


EM (EI, m/e): 442 (M^+), 444 ($\text{M} + 2^+$), 446 ($\text{M} + 4^+$)

EXAMPLE 2: Preparation of 3-nitrooxyethyl 2-(2,6-dichlorophenylamino) phenylacetoxypentanoate.

Solve 4.02 g (0.024 mol) of N,N'-carbonyl diimidazole in 30 ml of anhydrous chloroform under a dry nitrogen atmosphere.

Shake and add 8.78 g (0.024 mol) of 2-(2,6-dichlorophenylamino) phenylacetoxypentanoic acid in three shares. Shake for five minutes and add a solution of 3 g (0.024 mol) of 3-nitrooxypropanol to 15 ml of chloroform. Protect from the light and shake for 16 hours under these conditions. Eliminate the solvent by means of rotating steam and solve again in 100 ml of ethyl acetate. Wash three times with 50 ml of HCl 1N, twice with a solution of NaHCO_3 at 10% and neutralize by washing twice with water saturated with NaCl.

Dry on anhydrous sodium sulphate. Filter, eliminate the solvent at a reduced pressure to obtain an oil that crystallizes in methanol. Recover 5.99 g (54%) of 3-nitrooxypropyl 2-(2,6-dichlorophenylamino) phenylacetoxypentanoate (m.p. 57-59°C) with the following spectroscopic features:



IR (KBr, cm^{-1}): 3360 (N-H), 1755 (O-CH₂-CO-O), 1743 (ArCH₂-CO-O), 1629 (ONO₂).

¹H-RMN (DCCl₃, 8): 1.97 (m, 2H, J = 6.3 Hz, CH₂ (11)), 3.93 (s, 2H, CH₂ (8)), 4.22 (t, 2H, J_{10,11} = 6.1 Hz, CH₂ (10)), 4.41 (t, 2H, J_{12,11} = 6.1 Hz, CH₂ (12)), 4.69 (s, 2H, CH₂ (9)), 6.56 (d, 1H, J_{4,5} = 8.1 Hz, C-H (4)), 6.72 (s, 1H, N-H), 6.97 (m, 1H, J_{6,5} = J_{6,7} = 7.8 Hz, J_{6,4} = 0.9 Hz, C-H (6)), 6.99 (t, 1H, J_{1,2-3} = 8.1 Hz C-H (1)), 7.14 (m, 1H, J_{5,6} = J_{5,4} = 7.8 Hz, J_{5,7} = 1.8 Hz, C-H (5)), 7.27 (m, 1H, J_{7,6} = 7.8 Hz, J_{7,5} = 1.8 Hz C-H (7)), 7.34 (d, 2H, J_{2-3,1} = 8.4 Hz C-H (2) (3)).

¹³C-RMN (DCCl₃, 8): 26.12, 37.99, 61.15, 61.32, 69.37, 118.35, 122.13, 123.66, 124.16, 128.19, 128.86, 129.48, 130.93, 137.67, 142.67, 167.23, 171.42.

EM (EI, m/e): 456 (M)⁺, 458 (M + 2)⁺, 460 (M + 4)⁺.


EXAMPLE 3: Preparation of 4-chlorobutyl 2-(2,6-dichlorophenylamino) phenylacetoxycetate.

Solve 6.48 g (0.040 mol) of N,N'-carbonyl diimidazole in 100 ml of anhydrous dichloromethane under a dry nitrogen atmosphere. Shake and add 13.86 g (0.039 mol) of 2-(2,6-dichlorophenylamino)phenylacetoxycetic acid for five minutes in three shares.

Shake for five minutes and add a solution of 4.25 g (0.039 mol) of 4-chloro-1-butanol to 50 ml of dichloromethane. Protect from the light and shake for 20 hours.

Eliminate the solvent under reduced pressure and solve again the residue in 300 ml of ethyl acetate. Wash three times with 100 ml of HCl 1N, twice with a solution of NaHCO₃ at 10% and neutralize by washing twice with a saturated solution of NaCl. Dry on anhydrous sodium sulphate.

Filter, eliminate the solvent at a reduced pressure and purify the oil obtained by chromatography in silica gel column. Then eluting with dichloromethane to recover 11.95 g (69%) of 4-chlorobutyl 2-(2,6-dichlorophenylamino) phenylacetoxycetate in form of oil that has the following spectroscopic features:



IR (KBr, cm^{-1}): 3360 (N-H), 1742 (C=O).

^1H -RMN (DCCl_3 , 8): 1.75 (m, 4H, CH_2 (11)), CH_2 (12)), 3.50 (t, 2H, $J_{10,11} = 6$ Hz CH_2 (10)), 3.94 (s, 2H, CH_2 (8)), 4.17 (t, 2H, $J_{13,12} = 6$ Hz CH_2 (13)), 4.68 (s, 2H, CH_2 (9)), 6.56 (dd, 1H, $J_{4,5} = 7.8$ Hz, $J_{4,6} = 0.9$ Hz, C-H (4)), 6.74 (s, 1H, N-H), 6.98 (m, 1H, $J_{6,5} = J_{6,7} = 7.5$ Hz, $J_{6,4} = 1.2$ Hz, C-H (6)), 6.99 (t, 1H, $J_{1,2-3} = 7.9$ Hz, C-H (1)), 7.14 (m, 1H, $J_{5,6} = J_{5,4} = 7.4$ Hz, $J_{5,7} = 1.5$ Hz, C-H (5)), 7.27 (dd, 1H, $J_{7,6} = 7.5$ Hz, $J_{7,5} = 1.2$ Hz, C-H (7)), 7.34 (d, 2H, $J_{2-3,1} = 8.1$ Hz, C-H (2), C-H (3)).


^{13}C -RMN (DCCl_3 , 8): 25.64, 28.64, 37.92, 44.17, 61.11, 64.49, 118.30, 122.02, 123.97, 128.03, 128.73, 129.38, 130.82, 137.64, 142.59, 167.30, 171.31.

EM (EI, m/e): 443 (M^+), 445 ($\text{M} + 2$) $^+$, 447 ($\text{M} + 4$) $^+$.

Example 4: Preparation of 4-iodinebutyl 2-(2,6-dichlorophenylamino) phenylacetoxycetate.

Solve 8.84 g (0.019 mol) of 4-chlorobutyl 2-(2,6-dichlorophenylamino) phenylacetoxycetate in 100 ml of anhydrous acetone, add 5.85 g (0.038 mol) of sodium iodide and warm by reflux for 40 hours. Cool it, filter the sodium chloride formed and eliminate the solvent at a reduced pressure. Solve again the residuem with a mixture of 150 ml of dichloromethane and 50 ml of water. Decant the organic phase and dry on anhydrous sodium sulphate.

Filter and concentrate by means of vacuum to have 10.76 g of an oil. When chromatographing it in a silica gel column, eluting with dichloromethane you recover 8.76 g (82%) of 4-iodinebutyl 2-(2,6-dichlorophenylamino) phenylacetoxycetate in the form of an oil that has the following spectroscopic features:



IR (KBr, cm^{-1}): 3360 (N-H), 1746 (C=O).

¹H-RMN (DCCl₃, 8): 1.70 (m, 2H, CH₂, (11)), 1.80 (m, 2H, CH₂ (12)), 3.13 (t J_{10,11} = 6.6 Hz, 2H, CH₂ (10)), 3.93 (s, 2H, CH₂ (8)), 4.15 (t, J_{13,12} = 6.6 Hz, 2H, CH₂ (13)), 4.67 (s, 2H, CH₂ (9)), 6.56 (dd, 1H, J_{4,5} = 8.1 Hz, J_{4,6} = 1.2 Hz, C-H (4)), 6.73 (s, 1H, N-H), 6.98 (m, 1H, J_{6,5} = J_{6,7} = 7.3 Hz, J_{6,4} = 1.2 Hz, C-H (6)), 6.99 (t, 1H, J_{1,2-3} = 8.1 Hz, C-H (1)), 7.14 (m, 1H, J_{5,6} = J_{5,4} = 7.8 Hz, J_{5,7} = 1.5 Hz, C-H (5)), 7.26 (dd, 1H, J_{7,6} = 7.6 Hz, J_{7,5} = 1.5 Hz, C-H (7)), 7.34 (d, 2H, J_{2-3,1} = 7.8 Hz, C-H (2) (3)).

¹³C-RMN (DCCl₃, 8): 5.74, 29.26, 29.62, 38.06, 61.25, 64.25, 188.44, 122.17, 123.83, 124.11, 128.18, 128.87, 129.51, 130.96, 137.79, 142.72, 167.42, 171.43.


EM (EI, m/e): 535 (M)⁺, 537 (M + 2)⁺, 539 (M + 4)⁺.

Example 5: Preparation of 4-nitrooxybutyl 2-(2,6-dichlorophenylamine) phenylacetoxycetate.

A. Add for twenty minutes a solution of 8.35 g (0.015 mol) of 4-iodinebutyl 2-(2,6-dichlorophenylamine) phenylacetoxycetate in 30 ml of acetonitrile to a solution of 3.26 g (0.019 mol) of silver nitrate in 50 ml of anhydrous acetonitrile, and shake for 18 hours.

Finally, warm by reflux for 3 more hours. Filter and wash the precipitate with 30 ml of acetonitrile and concentrate the filtrates at a reduced pressure.

Shake the residue with 80 ml of ethyl ether and filter. Eliminate the ether and purify the oil obtained by silica gel chromatography. When eluting with dichloromethane you recover 6.48 g (88%) of 4-nitrooxybutyl 2-(2,6-dichlorophenylamine)phenylacetoxycetate inform of oil that has the following spectroscopic features:



IR (KBr, cm⁻¹): 3360 (N-H), 1743 (C = O), 1630 (ONO₂).

¹H-RMN (DDC1₃, 8): 1.69 (m, 4H, CH₂ (11) (12)), 3.93 (s, 2H, CH₂ (8)), 4.16 (t, 2H, J_{10,11} = 6 Hz, CH₂ (10)), 4.39 (t, 2H, J_{13,12} = 6 Hz, CH₂ (13)), 4.67 (s, 2H, CH₂ (9)), 5.55 (d, 1H, J_{4,5} = 7.8 Hz, C-H (4)), 6.73 (s, 1H, N-H), 6.97 (m, 1H, J_{6,5} = J_{6,7} = 7.5 Hz, J_{6,4} = 0.9 Hz, C-H (6)), 6.99 (t, 1H, J_{1,2-3} = 7.9 Hz, C-H (1)), 7.14 (m, 1H, J_{5,6} = J_{5,4} = 7.5 Hz, J_{5,7} = 1.5 Hz, C-H (5)), 7.26 (m, 1H, J_{7,6} = 6.9 Hz, J_{7,5} = 1.5 Hz, C-H (7)), 7.34 (D, 2H, J_{2-3,1} = 8.1 Hz, C-H (2) (3)).

¹³C-RMN (DCCl₃, 8): 23.35, 24.76, 38.01, 61.21, 64.42, 72.37, 118.39, 122.12, 123.76, 124.13, 128.16, 128.86, 129.49, 130.94, 137.73, 142.71, 167.36, 171.43.

EM (EI, m/me) 470 (M)⁺, 472 (M + 2)⁺, 474 (M + 4)⁺.


B. Add for twenty minutes a solution of 2.45 g (0.005 mol) of 4-chlorobutyl 2-(2,6-dichlorophenylamine) phenylacetoxycetate in 15 ml of acetonitrile to a solution of 1.5 g (0.008 mol) of silver nitrate in 20 ml of anhydrous acetonitrile, and warm by reflux for 12 hours. Add 1.5 g (0.008 mol) of silver nitrate and warm for 7 more hours. Filter the precipitate and wash it with 20 ml of acetonitrile and concentrate at a reduced pressure. Treat the residuum with 50 ml of ether and filter. Eliminate the ether by means of rotating steam to obtain an oil that is purified by chromatography in column on silica gel. When eluting with dichloromethane you recover 1.61 g (62%) of 4-nitrobutyl 2-(2,6-

dichlorophenylamino) phenylacetoxycetate in form of oil with spectroscopic features similar to the ones of the product obtained by method A.

Example 6: Preparation of N-(2-nitrooxyethyl)-2-(2,6-dichlorophenylamino) phenylacetoxycetamid.

Solve 6.86 g (0.042 mol) of N,N'-carbonyl diimidazole in 150 ml of anhydrous dichloromethane, in a dry nitrogen atmosphere. Shake and add 15 g (0.042 mol) of 2-(2,6-dichlorophenylamino) phenylacetoxycetic acid in four shares for five minutes. Shake for five minutes and add 7.16 g (0.042 mol) of 2-nitrooxyethylamine nitrate. Protect from the light and shake for 18 hours.

Eliminate the solvent by means of rotating steam and solve again the residue in 200 ml of ethyl acetate. Wash three times with 80 ml of HCl 1N, twice with NaHCO₃ at 10% and neutralize by washing twice with 80 ml of NaCl saturated solution. Dry on anhydrous sodium sulphate. Filter and eliminate the solvent at a reduced pressure to have an oil that crystallizes with toluene. Recover 6.60 g (35%) of N-(2-nitrooxyethyl)-2-(2,6-dichlorophenylamino) phenylacetoxycetamid in form of crystals (m.p. 87-88°C) with the following spectroscopic features:



IR (KBr, cm⁻¹): 3378 (N-H), 3298 (CON-H), 1751 (COO), 1677 (CONH), 1621 (ONO₂).


¹H-RMN (DCCl₃, 8): 3.48 (dt, 2H, J = 5.1 Hz, CH₂ (10)), 3.92 (s, 2H, CH₂ (8)), 4.42 (t, 2H, J = 5.1 Hz, CH₂ (11)), 4.65 (s, 2H, CH₂ (9)), 6.08 (s wide N-H), 6.56 (dd, 1H, J_{4,5} = 7.8 Hz, J_{4,6} = 0.9 Hz, C-H (4)), 7.00 (m, 1H, J_{6,5} = J_{6,7} = 7.5 Hz, J_{6,4} = 0.9 Hz C-H (6)), 7.02 (t, 1H, J_{1,2-3} = 7.8 Hz, C-H (1)), 7.17 (m, 1H, J_{5,6} = J_{5,4} = 7.5 Hz, J_{5,7} = 1.8 Hz, C-H (5)), 7.28 (dd, 1H, J_{7,6} = 7.4 Hz, J_{7,5} = 1.8 Hz, C-H (7)), 7.35 (d, 2H, J_{2,3,1} = 8.1 Hz, C-H (2) (3)).

¹³C-RMN (DCCl₃, 8): 36.54, 38.19, 63.05, 71.14, 118.22, 122.27, 123.12, 124.66, 128.57, 128.93, 129.77, 130.81, 137.24, 142.50, 167.34, 170.27.

EM (EI, m/e): 441 (M)⁺, 443 (M + 2)⁺.

Claims

1. Nitric ester of a 2-(2,6-dihalophenylamino) phenylacetoxyacetic acid derivative having the general formula (I):



(I)

where:


A is the fluorine, chlorine or bromine;

X means oxygen, NH or NR where R means a linear or ramified alkyl chain of 1 to 6 carbon atoms;

R₁ and R₂ mean, independently, hydrogen or a linear or ramified alkyl chain of 1 to 6 carbon atoms;

n is a number between 1 to 10.

2. Nitric ester of a 2-(2,6-dihalophenylamino) phenylacetoxyacetic acid derivative according to claim 1, where A is chlorine, X is oxygen, R₁ and R₂ are both hydrogen and n is 2.
3. Nitric ester of a 2-(2,6-dihalophenylamino) phenylacetoxyacetic acid derivative according to claim 1, where A is chlorine, X is oxygen, R₁ and R₂ are both hydrogen and n is 3.
4. Nitric ester of a 2-(2,6-dihalophenylamino) phenylacetoxyacetic acid derivative according to claim 1, where A is chlorine, X is oxygen, R₁ and R₂ are both hydrogen and n is 4.
5. Nitric ester of a 2-(2,6-dihalophenylamino) phenylacetoxyacetic acid derivative according to claim 1, where A is chlorine, X is NH, R₁ and R₂ are both hydrogen and n is 2.
6. Process of preparing nitric ester of 2-(2,6-dihalophenylamino) phenylacetoxyacetic acid derivatives with the general formula (I):




(I)

where:


A is fluorine, chlorine or bromine;
 X means O, NH or NR where R means a linear or ramified alkyl chain of 1 to 6 carbon atoms;
 R₁ and R₂ mean, independently, hydrogen or a linear or ramified alkyl chain of 1 to 6 carbon atoms;
 n is a number between 1 to 10,

comprising the step of condensing the 2-(2,6-dihalophenylamino) phenylacetoxymethyl ester with the formula (IV) where A is fluorine, chlorine or bromine



(IV)

with a derivative with the general formula (V)




(V)

where Y can be OH, NH₂ or NHR, where R is a linear or ramified alkyl chain of 1 to 6 carbon atoms, R₁ and R₂

mean, independently, hydrogen or a linear or ramified alkyl chain of 1 to 6 carbon atoms and n is a number between 1 to 10, in presence of an suitable condenser agent in an aprotic organic solvent.

7. Process according to claim 6 characterised in that the condenser agent is the N,N'-carbonyl diimidazol.
8. Process of preparing nitric esters of 2-(2,6-dihalophenylamino) phenylacetoxycetic acid derivatives having the general formula (I):




(I)

wherein:

- A is fluorine, chlorine or bromine;
 X means O, NH or NR where R means a linear or ramified alkyl chain of 1 to 6 carbon atoms;
 R₁ and R₂ mean, independently, hydrogen or a linear or ramified alkyl chain of 1 to 6 carbon atoms;
 n is a number between 1 to 10,

comprising the steps of:

- (i) condensation of 2-(2,6-dihalophenylamino) phenylacetoxycetic acid with the formula IV:




(IV)

where:


EP 0 738 706 A1

A is fluorine, chlorine or bromine with a general formula compound with the formula VI:



(VI)


where Y means OH, NH₂ or NHR where R, R₁, R₂ and n have the meaning above mentioned and Z is chlorine or bromine, in presence of a suitable condensation agent to provide the compound with the formula VII:



(VII)

where A, R₁, R₂ and Z have the meaning above mentioned and X means O, NH or NR being R a linear or ramified alkyl chain of 1 to 6 carbon atoms;

(ii) optional substitution of Z of the compound with the formula VII by a iodine atom by reaction with sodium iodide to provide the compound with the formula VIII:



(VIII)

iii) displacement of Z of the compound with the formula VII or, otherwise, of the iodine atom of the compound with the formula VIII with AgNO₃ in acetonitrile as solvent, to provide the corresponding nitroesters with the formula I.

9. Process according to claim 8 characterized in that the condensation agent in step (i) is the N,N'-carbonyl diimide.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 96 10 6009

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
A,D	EP-A-0 119 932 (PRODES S.A.) * abstract *	1	C07C229/42 C07C235/08 A61K31/215 A61K31/165
A,D	W0-A-94 04484 (CORLAY S.L.) * abstract *	1	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C07C A61K
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 12 July 1996	Examiner Rufet, J
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 01.92 (P04C01)